

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/316998981>

SOIL ENVIRONMENTAL METALS SPECIATION AND ASSOCIATED HEALTH RISKS IN SELECTED EDIBLE LEAFY PLANTS ON AMAÉCHI AND FOUR-CORNER DUMPSITES IN ENUGU OF ENUGU STATE, NIGERIA

Article · October 2014

CITATIONS

0

READS

44

5 authors, including:



Nwogo Ajuka Obasi

Alex Ekwueme Federal University Ndufu Alike Ikwo

84 PUBLICATIONS 1,023 CITATIONS

[SEE PROFILE](#)



Sunday Oge Elom

Federal University Ndufu Alike Ikwo

35 PUBLICATIONS 208 CITATIONS

[SEE PROFILE](#)



Stella Eberechukwu Obasi

Akanu Ibiam Federal Polytechnic, Unwana

7 PUBLICATIONS 12 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Physicochemical and Soil Enzymatic Activities as influenced by Mining operations in the Ishiagu area of Ebonyi State Nigeria [View project](#)



research article [View project](#)



International Journal of Life Sciences Biotechnology and Pharma Research





Research Paper

SOIL ENVIRONMENTAL METALS SPECIATION AND ASSOCIATED HEALTH RISKS IN SELECTED EDIBLE LEAFY PLANTS ON AMAÉCHI AND FOUR-CORNER DUMPSITES IN ENUGU OF ENUGU STATE, NIGERIA

N A Obasi^{1*}, S O Elom¹, C O Edeogu¹, C O Alisa² and S E Obasi³

*Corresponding Author: **N A Obasi** ✉ naobasi@yahoo.com

This work evaluated environmental metals speciation and the associated health risks using five selected edible leafy plants in Amaéchi and Four-Corner dumpsites in Enugu, Enugu State, Nigeria. The soil and plants samples were obtained from Amaéchi and Four-Corner dumpsites in Enugu and a nearby farm land (control site) and were subjected to standard methods of chemical analysis. Results obtained showed that mean soil physicochemical parameters and total extractable metals were significantly higher ($P < 0.05$) in the dumpsites compared to control site. Carbon:Nitrogen (C:N) ratio was significantly higher ($P < 0.05$) in the control site compared to that in Amaéchi dumpsite. Sequential extraction showed higher percentages (%) of the non-residual fraction for all the metals studied except Cu and Cr. The mean order of mobility and bioavailability of the metals were: $Pb > Fe > Cd > Zn > Mn > Cr > Ni > Cu$. Total mean concentration of metals in roots, stems and leaves of *Amaranthus hybridus*, *Talinum triangulare*, *Carica papaya*, *Ipomea batatas* and *Luffa aegyptica* were significantly higher ($P < 0.05$) in the dumpsites compared to control site. The translocation factor, biological concentration factor and biological accumulation coefficient values of the plant species varied for all the metals. The health risks implications of the study were discussed.

Keywords: Bioavailability, Phytoaccumulation, Dumpsites, Amaéchi, Four-Corner, Enugu

INTRODUCTION

Environmental metals are non-biodegradable and they are toxic to flora and fauna in the ecosystem when they exceed their threshold limit (Benjamin and Mwashote, 2003; Ikem *et al.*, 2003;

Krissanakriangkrai *et al.*, 2009; Ozturk *et al.*, 2009). Reports have shown that solid wastes dumps are sources of environmental pollution that introduce intolerable doses of heavy metals into surrounding soil and ground water (Uba *et al.*,

¹ Department of Medical Biochemistry, Federal University Ndufu-Alike, Ikwo-Nigeria.

² Department of Chemistry, Federal University of Technology, Owerri-Nigeria.

³ Department of Science Laboratory Technology, Akanu Ibiam Federal Polytechnic Unwana-Nigeria.

2008; Nubi *et al.*, 2008). Enugu is the state capital of Enugu State south-east, Nigeria with high population density, increased commercial and industrial activities and these have been associated with high refuse disposal rate (Ogwueleka, 2009; Abul, 2010).

Chemical fractionation of soil environmental metals shows that they are associated with organic matter, adsorbed onto Fe/Mn oxides or complexed with hydroxides, sulphides and carbonates (Tessier *et al.*, 1979; Kabata-Pendias, 2004). This speciation determines the relative mobility and bioavailability of the metals in the ecosystem and when the exchangeable, acid soluble and easily reducible fraction are high, the metal pollution is said to be of anthropogenic origin (Ewa-Szarek *et al.*, 2006; Obasi *et al.*, 2012). It is the chemical form or species of these heavy metals that is important in assessing their impacts on the environment because their chemical forms control their bioavailability or mobility which ultimately control heavy metal soil-plant transfer (Gupta and Sinha, 2006).

Dumpsites depending on age support flora and fauna due to the high organic manure contents which makes the soil fertile. Poor dumping of domestic, commercial and industrial refuse by most State Environmental Protection Agency in Nigeria on farmlands has led to the use of such dumpsites for cultivating varieties of edible vegetables and plant based foodstuff since in most cases, the soils are fertile and support plant growth. This practice gives serious health concern due to existing data on heavy metals phyto-accumulation potentials of plants in metal contaminated and polluted soils (Benson and Ebong, 2005). Reports have shown varying degree of phytotoxicity effects of these metals in

food chain and food web within the ecosystem (Ellis and Salt, 2003; Jarup, 2003).

Amaéchi and Four-Corner areas are suburb in Enugu metropolis where Enugu State Environmental Protection Agency dumps most of their evacuated refuse in their farmland. The preoccupation of the residents in these areas is farming and they extensively use these dumpsites as arable lands for cultivating varieties of edible vegetables and plant-based foodstuff. Since there is no proper routine assessment of the associated health risks, there may be danger along the food chain and food web at the future. This research therefore aims at providing baseline data that could be used to assess the futuristic health risks associated with such practice.

MATERIALS AND METHODS

Refuse Waste Soil Collection

Refuse waste soils were collected from two dumpsites, Amaéchi (Latitude 06 23' 08.32", Longitude 007 13' 38.50") and Four-Corner (Latitude 06 22' 09.68", Longitude 007 11' 18.45") and from the control site (Latitude 06 21' 58.55", Longitude 007 09' 36.25"), which is a farm land situated within the region. Triplicate sample from each dumpsite and control site were collected 7 m within the vicinity of the sites and composite samples were made in the laboratory. The samples were air dried, ground using manual soil grinder (DGSI Geotechnical instrumentation Model S-178), sieved (using 2 mm sieve), put in polythene bags and kept in glass desiccators (Baroda Scientific Glass Works) until analysis. During soil sample collection, care was taken to ensure that top soil at 0-20 cm depth from the rhizosphere of the studied plants were obtained from each site from where plant samples were rooted.

Dumpsite/Control Site Plant Sample Collection

Five dominant edible plant species within each study location: *Amaranthus hybridus*, *Talinum triangulare*, *Carica papaya*, *Ipomea batatas* and *Luffa aegyptia* were obtained and used for the study. A total of 6-10 plant samples of each plant species were randomly uprooted and collected from each of the dumpsite and control site and separately mixed to form a composite sample, placed in labeled pre-cleaned polythene bags and transported within 14 h to the Chemistry Laboratory of National Research Institute for Chemical Technology, Zaria, Nigeria for further analysis. Before analysis, plant roots and a mixture of the stems and leaves (shoots) were carefully removed and washed (for 2-3 min approximately) with tap water and deionized water to remove any soil and surface dust. Plant samples were dried at room temperature for a day, oven dried at 80°C to constant weight and pulverized to fine powder using milling grinder (Thomas Wiley Model 4). Ground plant samples collected in labeled pre-cleaned polythene bags were stored in glass desiccators (Baroda Scientific Glass Works).

Physicochemical Analysis of Samples

Soil pH was determined using digital pH meter (Jenway 3015) at a ratio of 1:2.5 soil/water according to the procedure described by Bates (1954). Soil electrical conductivity was determined using digital electrical conductivity meter (Jenway 615D) according to the procedure outlined by Whitney (1998) with some modifications. The soil moisture content was determined according to the procedure outlined by APHA (1998) while the cation exchange capacity of the soil samples were determined by

ammonium saturation method described by Dewis and Freitas (1970). Organic carbon and organic matter were determined according to the procedure outlined by Osuji and Adesiyan (2005) while total nitrogen was determined as described by Yeomans and Bremmer (1991). SO_4^{2-} was quantified by the procedure described by Butters and Chenery (1959) and PO_4^{3-} was determined by procedure described by Olsen and Sommers (1982), respectively.

Sequential Extraction of Heavy Metals

The conventional method developed by Tessier *et al.* (1979) as outlined with modifications in Obasi (2012) was employed for the sequential extraction of heavy metals.

Determination of Heavy Metals in Plant Species

The mineral elements comprising cadmium (Cd), copper (Cu), manganese (Mn), lead (Pb), zinc (Zn), iron (Fe), nickel (Ni) and chromium (Cr) were determined according to the procedure described by Obasi (2012) using atomic absorption spectrophotometer (Bulk Scientific Model 210 VGP).

Determination of Phytoremediation Quotient

The Translocation Factor (TF) defined as the ratio of heavy metals in plant shoot to that in plant root was calculated using the procedure described by Cui *et al.* (2007).

$$\text{i.e. Translocation factor (TF)} = \frac{[\text{Metals}]_{\text{shoot}}}{[\text{Metals}]_{\text{root}}}$$

The Biological Concentration Factor (BCF) was calculated as metal concentration ratio of plant roots to soil as described by Yoon *et al.* (2006).

$$\text{i.e. Biological Concentration Factor (BCF)} =$$

$$\frac{[\text{Metals}]_{\text{root}}}{[\text{Metals}]_{\text{soil}}}$$

Biological Accumulation Coefficient (BAC) was calculated as a ratio of heavy metal in shoots to that in soil as described in the procedure by Li *et al.* (2007).

i.e. Biological Accumulation Coefficient (BAC) = $\frac{[\text{Metals}]_{\text{shoot}}}{[\text{Metals}]_{\text{soil}}}$

STATISTICAL ANALYSIS

The experimental results were expressed as mean \pm standard deviation (SD) of triplicate determinations. Analysis of variance for all the measured variables was performed by SPSS version 9.2 (Inc., Chicago, USA) software and significant differences were shown at $P < 0.05$ (Kerr *et al.*, 2002).

RESULTS

The results of soil physico-chemical properties are shown in Table 1. Results obtained showed

that mean pH, electrical conductivity, moisture, cation exchange capacity, total organic carbon, total organic matter, total nitrogen, phosphate, and sulphate, were significantly higher ($P < 0.05$) in the dumpsites compared to control site while carbon:nitrogen ratio in Amaéchi was significantly lower ($P < 0.05$) compared to control site. The results of the sequential extractions of the heavy metals are shown in Tables 2a and 2b. The results indicated that total extractable metals were significantly ($P < 0.05$) higher in all the dumpsites compared to the control site. In all cases, the highest total extractable metals were present in Amaéchi dumpsite followed by Four-Corner dumpsite except for total extractable Cu where the reverse was the order. Higher percentages (%) of the non-residual fraction were observed for all the metals studied except Cu and Cr. The mean percentage order of mobility and bioavailability of these metals (Tables 2a and 2b) were: Pb > Fe > Cd > Zn > Mn > Cr > Ni > Cu.

Table 1: Physico-chemical Parameters of Soils in Studied Dumpsites

Sites/Parameter	A	B	AB
pH(H ₂ O)	7.90 \pm 0.00 ^c	7.65 \pm 0.01 ^b	7.10 \pm 0.02 ^a
Electrical Conductivity (mScm-1)	2.78 \pm 0.02 ^c	1.96 \pm 0.02 ^b	0.95 \pm 0.01 ^a
Moisture (%)	39.60 \pm 0.04 ^a	43.00 \pm 0.02 ^b	40.15 \pm 0.03 ^a
Cation Exchange Capacity (Cmol/kg)	15.10 \pm 0.00 ^b	18.40 \pm 0.03 ^c	10.20 \pm 0.02 ^a
Total Organic Carbon (%)	3.00 \pm 0.00 ^b	3.18 \pm 0.02 ^b	1.02 \pm 0.02 ^a
Total Organic Matter (%)	5.17 \pm 0.09 ^b	5.48 \pm 0.02 ^c	1.78 \pm 0.02 ^a
Total Nitrogen (%)	0.28 \pm 0.02 ^b	0.24 \pm 0.02 ^a	0.25 \pm 0.01 ^a
PO ₄ ³⁻ (%)	186.70 \pm 0.02 ^c	165.50 \pm 0.04 ^b	142.40 \pm 0.10 ^a
SO ₄ ²⁻ (%)	13.59 \pm 0.13 ^c	12.00 \pm 0.05 ^a	12.26 \pm 0.02 ^{ab}
C:N RATIO	10.71 ^a	13.25 ^c	11.52 ^b

Note: Values are mean of three (n=3) replicates \pm standard deviation; A = Amaéchi dumpsite, B = Four-Corner dumpsite, AB = Control site; Figures followed by the same alphabets along the row are not significantly different at $P < 0.05$ using Duncan Multiple Range Test (DMRT).

The results of total heavy metals concentration (mg/kg) in roots and shoots of plant species are shown in Tables 3a and 3b. Total mean concentration of metals in different parts of *Amaranthus hybridus*, *Talinum triangulare*, *Carica papaya*, *Ipomea batatas* and *Luffa aegyptica* were significantly higher ($P < 0.05$) in the dumpsites compared to control site. The results also showed that different plant species absorbed metals at varying concentrations in their various parts (Tables 3a and 3b). The results (Figure 1) indicated that Translocation Factor (TF) values vary from one plant species to another and from one heavy metal to another. The results indicated that *A. hybridus* had TF > 1 for Mn, Zn,

Fe and Cr, *T. triangulare* had TF > 1 for Zn, Fe and Cr, *C. papaya* had TF > 1 for all the studied metals except Ni and Pb while *I. batatas* and *L. aegyptica* had TF > 1 for Cd, Cu, Mn, Pb, Zn, Fe, Ni and Cr (Figure 1).

Figure 2 shows the results of BCF of the five plant species for the different metals. The results (Figure 2) showed that only *A. hybridus* and *C. papaya* had BCF > 1 for Cu and Fe respectively. The results of BAC are shown in Figure 3. The results (Figure 3) showed that *C. papaya* had BAC > 1 for Cu and Fe in all the sites while *A. hybridus*, *T. triangulare* and *L. aegyptica* had BAC > 1 for Fe in Amaéchi dumpsite only.

Table 2a: Heavy Metal Concentrations (Mg/Kg) in Each Fraction of Waste Soils in Studied Dumpsites

Sites/Fractions	Cd			Cu			Mn			Pb		
	A	B	AB	A	B	AB	A	B	AB	A	B	AB
Exchangeable	9.63 ±0.05	Nd	0.98 ±0.01	Nd	2.10 ±0.02	Nd	7.26 ±0.02	Nd	0.41± 0.03	88.30± 0.02	12.95± 0.03	1.95± 0.02
Oxidizable	9.61 ±0.08	3.02 ±0.03	0.64 ±0.02	4.02 ±0.03	Nd	0.74 ±0.02	73.58 ±0.02	11.68 ±0.02	3.15 ±0.05	187.65 ±0.05	Nd	3.92 ±0.02
Acid Soluble	6.18 ±0.04	5.82 ±0.03	Nd	Nd	Nd	Nd	13.57 ±0.03	26.13 ±0.11	0.85 ±0.05	102.74 ±0.02	13.06 ±0.02	2.00 ±0.01
Reducible	9.63 ±0.11	7.04 ±0.02	1.05 ±0.05	Nd	Nd	0.26 ±0.02	96.05 ±0.04	108.34 ±0.16	3.40 ±0.02	37.44 ±0.02	Nd	3.81 ±0.03
Residual	11.70 ±0.02	10.06 ±0.03	1.12 ±0.02	9.44 ±0.02	13.18 ±0.04	2.10 ±0.05	7.20 ±0.02	18.61 ±0.07	1.07 ±0.03	202.70 ±0.05	13.48 ±0.02	3.24 ±0.02
Total Extractable Metals	46.75 ^c ±0.05	25.94 ^b ±0.04	3.79 ^a ±0.03	13.46 ^b ±0.02	15.28 ^c ±0.02	3.10 ^a ±0.00	198.26 ^c ±0.04	164.76 ^b ±0.12	8.88 ^a ±0.05	618.83 ^c ±0.11	39.49 ^b ±0.07	15.42 ^a ±0.02
Non-residual (%)	74.97	61.22	70.45	29.87	13.74	32.26	96.37	88.70	87.95	67.24	65.86	78.99
Residual (%)	25.03	38.78	29.55	70.13	86.26	67.74	3.63	11.30	12.05	32.76	34.14	21.01
Mobile Phase (%)	33.82	22.44	25.86	0.00	13.74	0.00	10.51	15.86	14.19	30.87	65.86	25.62

Note: Values are mean of three (n=3) replicates ± standard deviation; A = Amaéchi dumpsite, B = Four-Corner dumpsite, AB = Control site; Figures followed by the same alphabets along the row are not significantly different at $P < 0.05$ using Duncan Multiple Range Test (DMRT).

**Table 2b: Heavy Metal Concentrations (Mg/Kg)
in Each Fraction of Waste Soils in Studied Dumpsites Continued**

Sites/Fractions	Cd			Cu			Mn			Pb		
	A	B	AB	A	B	AB	A	B	AB	A	B	AB
Exchangeable	37.15 ±0.03	Nd	1.46 ±0.02	35.88 ±0.02	Nd	11.2 ±0.02	Nd	Nd	0.2 ±0.01	2.9 ±0.02	Nd	0.22 ±0.02
Oxidizable	48.5 ±0.02	34.4 ±0.00	2.96 ±0.03	23.11 ±0.07	19.2 ±0.02	22.05 ±0.03	8 ±0.00	8.4 ±0.02	0.95 ±0.01	Nd	3.14 ±0.03	0.4 ±0.02
Acid Soluble	13.14 ±0.02	26.31 ±0.05	1 ±0.05	31.45 ±0.03	21.43 ±0.05	10.22 ±0.06	13.95 ±0.03	Nd	Nd	2.85 ±0.02	Nd	Nd
Reducible	42.36 ±0.03	25.38 ±0.07	2.64 ±0.02	26.18 ±0.02	27 ±0.01	21.11 ±0.03	Nd	5.4 ±0.02	0.8 ±0.02	Nd	2.22 ±0.05	0.34 ±0.02
Residual	79.8 ±0.02	100.29 ±0.03	3.08 ±0.02	37.7 ±0.02	37.28 ±0.12	24.82 ±0.16	26.66 ±0.08	7.85 ±0.03	1.35 ±0.02	16.05 ±0.04	13 ±0.02	0.81 ±0.03
Total Extractable Metals	220.95 ^c ±0.05	186.38 ^b ±0.02	11.14 ^a ±0.02	154.32 ^c ±0.11	104.91 ^b ±0.09	89.40 ^a ±0.02	48.61 ^c ±0.17	21.65 ^b ±0.08	3.30 ^a ±0.02	21.80 ^c ±0.02	18.36 ^b ±0.04	1.87 ^a ±0.03
Non-residual (%)	63.88	46.19	72.35	75.57	64.46	72.24	45.16	63.74	50.09	26.38	29.19	56.68
Residual (%)	36.12	53.81	27.65	24.43	35.54	27.76	54.84	36.26	40.91	73.62	70.81	43.32
Mobile Phase (%)	22.76	14.12	22.08	43.63	20.43	23.96	28.7	0	6.06	26.38	0	11.76

Note: Values are mean of three (n=3) replicates ± standard deviation; A = Amaéchi dumpsite, B = Four-Corner dumpsite, AB = Control site; Figures followed by the same alphabets along the row are not significantly different at P < 0.05 using Duncan Multiple Range Test (DMRT).

DISCUSSION

The observed mean pH shows that the dumpsites soils are alkaline which is in line with earlier reports on dumpsites (Uba *et al.*, 2008; Obasi, 2012; Obasi *et al.*, 2012). The observed high conductivity value of the waste soil is an indication that there are high soluble salts in the soil and this may be due to the presence of metal scraps in the refuse dumpsite (Karaca, 2004; Arias *et al.*, 2005; Obasi 2012). The moisture content was observed to be generally high and it is expected considering the overall climatic condition of the area under study. Cation exchange capacity observed in the studied sites fall within permissible range for agricultural lands and may have impacted positively to the fertility of the soil (Yoo

and James, 2002). The observed mean percentages of Total Organic Carbon (TOC) and Total Organic Matter (TOM) values were high (Enwezor *et al.*, 1988) and thus may serve as an important indicator of the soil as a rooting environment (Okalebo *et al.*, 1993). The high concentration of the total nitrogen, PO_4^{3-} and SO_4^{2-} in the refuse waste soils and the high ratio of carbon to nitrogen (C:N) may have contributed to overall fertility of the soils which is an indication that the soils would support plant species diversity and growth (Okalebo, 1993; Obute *et al.*, 2010).

The observed high values of total extractable metals may be attributed to dumping of numerous metal containing wastes such as cadmium and lead acid batteries, metal scraps among others

in the dumpsites. These observed values were however below the permissible limits allowable for agricultural lands except for Cd (CCME, 1991; MAFF, 1992; CEC, 1986; USEPA, 1986). The high percentage of Cd in the mobile fractions suggests that Cd in these soils was potentially more bioavailable for plants uptake (Kuo *et al.*, 1983; Gupta and Sinha, 2006). Cu was mostly found in the residual phase (i.e., bound to silicates and detrital materials) which implicates association with the oxidizable fraction (bound to organic matter) as in organic copper complexes (Stumm and Morgan, 1981). The high percentage of Mn in the reducible phase in all the waste soils may be due to the precipitation of amorphous hydrous oxides of manganese as the dumpsites ages (Staelens *et al.*, 2000). High percentage of Pb,

Zn and Fe in the mobile phase (exchangeable and acid soluble phases) indicates high bio-availability and higher risks to the ecosystem (Kuo *et al.*, 1983). Most of the Ni and Cr were found in the residual and oxidizable fractions and this may be attributed to the alkaline stabilization process of the soils leading to organic complexation that impair their mobility (Tokalioglu *et al.*, 2000; Alvarez *et al.*, 2002).

The observed variation in the amount of metals accumulated by the different plant species in their various parts in the various sites is an indication that uptake of an element by a plant is primarily dependent on the plant species, its inherent controls, soil quality and metal concentrations in their habitual soil environment (Chunilall *et al.*, 2005). Thus, plant species significantly influence

Table 3a: Total Heavy Metals Concentration (Mg/Kg) In Roots and Shoots of Plant Species in the Studied Sites

Plant Species	<i>Amarathushybridus</i>		<i>Talinumtriangulare</i>		<i>Caricapapaya</i>		<i>Ipomeabatatas</i>		<i>Luffaegyptiaca</i>		
	Sites	Roots	Shoots	Roots	Shoots	Roots	Shoots	Roots	Shoots	Roots	Shoots
Cd	A	23.25±0.08	12.87±0.26	28.58±0.15	10.02±0.06	27.05±0.01	33.28±0.04	3.53±0.13	6.23±0.02	4.15±0.02	10.84±0.20
	B	19.44±0.02	7.95±0.02	19.45±0.01	16.80±0.01	16.93±0.02	17.24±0.03	2.08±0.01	3.04±0.02	2.46±0.02	6.55±0.03
	AB	2.44±0.55	1.26±0.02	2.31±0.31	0.86±0.01	1.78±0.02	2.62±0.06	0.27±0.03	0.43±0.01	0.34±0.01	0.83±0.11
Cu	A	13.65±0.02	9.14±0.02	7.25±0.04	4.16±0.01	6.60±0.10	15.08±0.13	3.42±0.02	5.40±0.10	2.87±0.05	4.62±0.02
	B	16.51±0.06	10.33±0.01	8.90±0.02	6.03±0.11	9.05±0.02	17.50±0.02	3.95±0.04	6.86±0.02	3.56±0.02	5.91±0.17
	AB	3.57±0.01	2.01±0.07	3.34±0.02	2.04±0.01	2.52±0.02	4.88±0.13	0.86±0.02	1.55±0.04	0.75±0.02	1.23±0.01
Mn	A	3.96±0.02	8.54±0.05	11.26±0.02	8.35±0.03	7.05±0.01	11.60±0.02	1.36±0.05	4.87±0.06	4.12±0.03	13.57±0.02
	B	3.39±0.03	7.48±0.02	8.50±0.01	4.77±0.07	8.67±0.03	14.59±0.01	1.30±0.02	4.19±0.01	3.85±0.04	8.96±0.02
	AB	0.53±0.05	0.66±0.02	1.21±0.03	0.73±0.01	4.16±0.02	7.75±0.02	0.32±0.01	1.18±0.01	2.46±0.02	6.15±0.04
Pb	A	10.80±0.02	6.95±0.04	7.75±0.04	4.96±0.02	15.14±0.03	11.68±0.02	4.37±0.03	6.60±0.01	5.14±0.02	7.65±0.04
	B	16.35±0.04	12.33±0.01	13.61±0.03	10.09±0.04	20.44±0.02	13.75±0.04	5.26±0.02	7.40±0.02	5.50±0.00	8.10±0.12
	AB	1.61±0.03	0.91±0.01	1.15±0.04	0.88±0.02	1.94±0.02	1.08±0.05	0.93±0.02	1.45±0.04	1.03±0.01	1.88±0.02

Note: Values are mean of three (n=3) replicates ± standard deviation; A = Amaéchi dumpsite, B = Four-Corner dumpsite, AB = Control site

Table 3b: Total Heavy Metals Concentration (Mg/Kg) In Roots and Shoots of Plant Species in the Studied Sites Continued

Plant Species	<i>Amarathushybridus</i>		<i>Talinumtriangulare</i>		<i>Caricapapaya</i>		<i>Ipomeabatatas</i>		<i>Luffaegyptiaca</i>		
	Sites	Roots	Shoots	Roots	Shoots	Roots	Shoots	Roots	Shoots	Roots	Shoots
Zn	A	24.95±0.04	27.92±0.02	22.36±0.04	25.05±0.03	26.72±0.02	33.52±0.02	14.97±0.03	18.35±0.04	17.85±0.03	20.17±0.09
	B	10.63±0.05	14.40±0.02	9.70±0.05	11.92±0.02	13.80±0.01	16.75±0.03	8.76±0.02	11.24±0.05	9.30±0.15	13.85±0.03
	AB	0.89±0.03	1.20±0.04	1.04±0.01	1.24±0.02	1.33±0.01	1.85±0.04	1.13±0.01	1.52±0.02	1.48±0.02	1.22±0.004
Fe	A	154.90±0.10	166.15±0.33	158.75±0.04	165.99±0.18	195.16±0.14	209.10±0.00	131.65±0.17	149.14±0.16	147.86±0.22	169.65±0.13
	B	88.61±0.13	92.33±0.21	86.41±0.16	89.48±0.07	107.82±0.14	123.66±0.18	87.16±0.05	88.06±0.02	83.08±0.23	94.84±0.15
	AB	59.33±0.15	63.96±0.04	60.67±0.23	64.09±0.16	87.11±0.09	90.27±0.29	58.36±0.23	61.93±0.15	59.93±0.17	62.98±0.14
Ni	A	3.46±0.02	3.12±0.02	2.98±0.05	2.66±0.02	4.57±0.03	4.35±0.04	2.20±0.02	2.35±0.04	2.32±0.02	2.46±0.02
	B	2.75±0.03	2.53±0.05	2.20±0.20	1.97±0.01	3.07±0.03	2.95±0.04	2.15±0.02	2.28±0.02	2.22±0.01	2.48±0.02
	AB	0.97±0.03	0.85±0.04	0.83±0.01	0.71±0.03	1.03±0.01	0.92±0.02	0.73±0.01	0.77±0.05	0.75±0.03	0.82±0.02
Cr	A	7.85±0.04	19.12±0.15	13.75±0.07	16.33±0.08	14.92±0.02	18.74±0.12	8.72±0.06	12.81±0.09	10.95±0.04	15.10±0.02
	B	5.62±0.02	7.19±0.14	4.96±0.06	6.88±0.02	6.86±0.04	9.39±0.13	3.11±0.03	5.99±0.07	6.33±0.05	7.40±0.02
	AB	0.65±0.01	0.93±0.05	0.89±0.01	1.08±0.02	1.01±0.03	1.26±0.02	0.84±0.02	0.98±0.04	0.76±0.02	0.87±0.03

Note: Values are mean of three (n=3) replicates ± standard deviation; A = Amaéchi dumpsite, B = Four-Corner dumpsite, AB = Control site

Figure 1: Translocation Factor (Tf) of Plants For All The Metals in the Studied Sites




Figure 2: Biological Concentration Factor (Bcf) Of Plants For All The Metals In The Studied Sites





Figure 3: Biological Accumulation Coefficient (Bac) Of Plants For All The Metals In The Studied Sites



their rate of metal uptake, storage and distribution to various parts and this may be due to their genetic variability (Shauibu and Ayodele, 2002; Ebong *et al.*, 2008). Since the rate of metal uptake is greatly influenced by plant species, the transfer factors of the metals by each plant species are desirable for classification of the plants' phytoaccumulation, photostabilization and phytoextraction potentials (Ayari *et al.*, 2010; Malik *et al.*, 2010).

The observed high level of metals in the different plant species grown on the dumpsites relative to those on the control sites is an indication that the level of metal in the soil-plants habitual environment influences its uptake (Zhu *et al.*, 1990; Ghosh and Singh, 2005). The ability of these plant species to accumulate the observed high level of metals may be due to the ability of the plants to evolve mechanisms that could enhance its phyto-accumulation potentials and metal detoxification. Most of the plant species accumulated these metals in relatively high amount that could be hazardous along food chain and food web of their dependent species (Vecera *et al.*, 1999; Assuncao *et al.*, 2003). However, this could only arise on continual dependence on plants grown on such contaminated environment for a long period of time since in some cases the level in the dumpsites does not considerably exceed those in the control site which compare favorably with established critical permissible limits in plants (Shanker *et al.*, 2005; Alloway, 1996).

A critical value greater than one (>1) for plants' Translocation Factors (TF), BCF and BAC is used to evaluate the potentials of plant species for phytoextraction, phytostabilization and phyto-remediation respectively (Yoon *et al.*, 2006; Cui

et al., 2007; Li *et al.*, 2007). High root to shoot translocation ($TF > 1$) as observed for some of the metals in some of the studied plants is an indication that some of these plants have vital characteristics to be used in phyto-extraction (Ghosh and Singh, 2005; Malik *et al.*, 2010). The observed high metal translocation quotient may be attributed to well-developed metal detoxification mechanism based on sequestration in the tolerant plant species (Ghosh and Singh, 2005; Cui *et al.*, 2007). The suitability of plant species with high TF values for phyto-extraction is due to their ability to translocate heavy metals to easily harvestable parts (shoots) (Yoon *et al.*, 2006; Cui *et al.*, 2007; Li *et al.*, 2007). Elevated concentration of heavy metals in roots of plants species and low translocation into above ground parts (BCF) indicate their suitability for phyto-stabilization (Ghosh and Singh, 2005). The observed plant species with $BCF > 1$ and $TF < 1$ may be useful for phyto-stabilization of one, two or more of the metal contaminants of the study area. Plants with well-developed cellular mechanisms for heavy metal detoxification and tolerance ($BAC > 1$) is used as an indicator of high heavy metals accumulator plant species (Ghosh and Singh, 2005) and when the plant is able to accumulate up to 1000 mg/kg of metal and above, the plant is classified as a hyper-accumulator (Baker and Brooks, 1989). The observed BAC values showed that the plants exhibited varying levels of phyto-accumulation potentials. However, none of the edible plants studied is a hyper-accumulator of any of the metals based on the critical set limit (Baker and Brooks, 1989).

CONCLUSION

The metal pollution index and health risks associated with Amaéchi and Four-Corner

dumpsites in Enugu, Enugu State, South-East, Nigeria have been highlighted. The dumpsites were observed to be fertile with higher percentage of non-residual fraction for all the metals studied except for Cu and Cr that had more abundant pool of residual fractions. Generally, the order of mobility and bioavailability of the metals were: Pb > Fe > Cd > Zn > Mn > Cr > Ni > Cu. Mean total extractable metals in the dumpsites relative to that in the control site showed pollution from anthropogenic sources. The edible plants dominant in the area (*A. hybridus*, *T. triangulare*, *C. papaya*, *I. batatas* and *L. aegyptiaca*) accumulated metals in varying degrees in their various parts and exhibited phyto-extraction, phyto-stabilization and phyto-accumulation potentials which are indicators of health risks along food chain and food web. The nutritional quality and long term effects of dependence on edible dumpsite plants as sources of vegetables need further investigation.


REFERENCES

1. Abul S (2010), "Environmental and health impact of solid waste disposal at Mangwaneni dumpsite in Manzini: Swaziland", *Journal of Sustainable Development in Africa*, Vol. 12, No. 7, pp. 64-73.
2. Alloway B J (1996), *Heavy metals in soils*, London: John Wiley and Sons Inc.
3. Alvarez E A, Mochon M C, Sanchez J C J and Rodriguez M T (2002), "Heavy metal extractable forms in sludge from wastewater treatment plants", *Chemosphere*, Vol. 47, pp. 765-775.
4. American Public Health Association. Standard methods of examination of water and waste water. USA:Washington, DC, 1998.
5. Arias M E, Gonzalez-Perez J A, Gonzalez-Villa F J and Ball A S (2005), "Soil health: A new challenge for microbiologists and chemists", *International Microbiology*, Vol. 8, pp. 13-21.
6. Assuncao A G L, Schat H and Aarts M G M (2003), "*Thlaspi caerulescens*, an attractive model species to study heavy metal hyperaccumulation in plants", *New Phytol.*, Vol. 159, pp. 351-360.
7. Ayari F, Hamdi H, Jedidi N, Gharbi N and Kossai R (2010), "Heavy metal distribution in soil and plant in municipal solid waste compost amended plots", *International Journal Environment, Science and Technology*, Vol. 7, No. 3, pp. 465-472.
8. Bates R G (1954), *Electromeric pH determination*, New York: John Willey and Sons Inc.
9. Baker A J M and Brooks R R (1989), *Terrestrial higher plants which hyperaccumulative metals*, New York: CAB International.
10. Benjamin M and Mwashot M (2003), "Levels of caesium and lead in water, sediment and selected fish species in Mombasa Kenya Western Indian", *Oceanic J. Mar. Sci.*, Vol. 2, pp. 25-34.
11. Benson N U and Ebong G A (2005), "Heavy metals in vegetables commonly grown in a tropical garden ultisol", *Journal of Sustainable Tropical Agricultural Resources*, Vol. 16, pp. 77-80.
12. Butters B and Chenery E M (1959),

- “Determination of sulphate in soil, plant materials and water by the turbidimetric method”, *Analyst Lond.*, Vol. 84, pp. 239-242.
13. Canadian Council of Ministers of the Environment (CCME) (1991), Interim Canadian environment quality criteria for contaminated sites. Canada: Report CCME EPC-CS3.
14. Council of the European Communities (CEC) (1986), “Directive of 12th June, 1986 on the protection of the environment and in particular soil when sewage sludge is used in agriculture. Official”, *Journal European Community*, L181:6-12.
15. Chunilall V, Kindness A and Johnalagada S B (2005), “Heavy metal uptake by two edible *Amaranthus* herbs grown on soils contaminated with lead, mercury, cadmium, and nickel”, *Journal of Environmental Science and Health*, Vol. 40, pp. 375-385.
16. Cui S, Zhou Q and Chao L (2007), “Potential hyper-accumulation of Pb, Zn, Cu and Cd in enduring plants distributed in an old semetry, northeast, China”, *Environmental Geology*, Vol. 51, pp. 1043-1045.
17. Dewis J and Freitas F (1970), *Physical and chemical methods of soil and water analysis* (Soil Bulletin 10) Rome: FAO.
18. Ebong G A, Akpan M M and Mkpenie V N (2008), “Heavy metal contents of municipal and rural dumpsite soils and rate of accumulation by *Carica papaya* and *Talinum triangulare* in Uyo, Nigeria”, *E-Journal of chemistry*, Vol. 5, No. 2, pp. 281-290.
19. Ellis D R and Salt D E (2003), “Plants, selenium and human health”, *Current Opinion Plant Biology*, Vol. 6, pp. 273-279.
20. Enwezor W O, Ohiri A C, Opubaribo E E and Udoh E J (1988), “A review of soil fertility investigators in south eastern Nigeria”, Nigeria:HFDA, Lagos.
21. Ewa-Szarek G, Amrowiez A and Gwazda R (2006), “Trace elements concentration in fish and bottom sediments of autotrophic dam reservoir”, *Int. J. Hydrol.*, Vol. 35, pp. 331-352.
22. Ghosh M and Singh S P (2005), “A review of phytoremediation of heavy metals and utilization of it’s by-products”, *Applied Ecology and Environmental Research*, Vol. 3, No. 1, pp. 1-18.
23. Gupta A K and Sinha S (2006), “Chemical fractionation and heavy metal accumulation in the plant of *Sesamum indicum* (L.) Var. T55 grown on soil amended with tannery sludge: selection of single extractants”, *Chemosphere*, Vol. 64, pp. 161-173.
24. Ikem A, Egiebog N O and Nyavor K (2003), “Trace Elements in water, fish and sediment from Tuskegee Lake, Southeastern USA”, *Water, Air Soil Pollut.*, Vol. 149, pp. 51-75.
25. Jarup L (2003), “Hazards of heavy metals contamination”, *British Medical Bulletin*, Vol. 68, pp. 167-182.
26. Kabata-Pendias A (2004), “Soil-plant transfer of trace elements: an environment issue”, *Geoderma*, Vol. 122, pp. 143-149.
27. Karaca A (2004), “Effect of organic wastes on the extractability of Cadmium, Copper, nickel and Zinc in Soil”, *Geoderma*, Vol. 122, pp. 297- 303.

28. Kerr A W, Hall H K and Kozub S A (2002), *Doing Statistics with SPSS*, London: SAGE Publications Ltd.
29. Krissanakriangkrai O, Suparpacboon W, Juwa S, Chacwong S and Swaddiwudhipong W (2009), "Bioavailable cadmium in water, sediment and fish, in a highly contaminated area of Thai-Myammy borde", *Thammasat Int. J. Sci. Technol.*, Vol. 14:60-68.
30. Kuo S, Heilman P E and Baker A S (1983), "Distribution and forms of Cu, Zn, Cd, Fe and Mn in soils near a Copper smelter", *Soil Sci.*, Vol. 135, pp. 101-109.
31. Li M S, Luo Y P and Su Z Y (2007), "Heavy metals concentrations in soils and plant accumulation in a restored manganese mine land in Guangxi, South China", *Environmental pollution*, Vol. 147, pp. 168-175.
32. Ministry of Agricultural, Forestry and Fisheries (MAFF) (1992), "Code of good agricultural practice for the protection of soil", Welch Office Agriculture Department, Draft Consultation Document. London: Ministry of Agricultural, Forestry and Fisheries.
33. Malik R N, Husain S Z and Nazir I (2010), "Heavy metal contamination and accumulation in soil and wild plant species from industrial area of Islamabad, Pakistan", *Pakistan Journal Botany*, Vol. 42, No. 1, pp. 291-301.
34. Nubi O A, Osibanjo O and Nubi A T (2009), "Impact assessment of dumpsite leachate on the qualities of surface water and sediment of River Eku, Ona-ara Local Government, Oyo State, Nigeria", *Science World Journal*, Vol. 3, No. 3, pp. 17-20.
35. Obasi N A, Akubugwo E I, Ugboogu O C and Chinyere G C (2012), "Heavy Metals Bioavailability and Phyto-accumulation Potentials of Selected Plants on Burrow-pit Dumpsites in Aba and Ntigha Dumpsite in Isiala Ngwa of Abia State, Nigeria", *Nigerian Journal of Biochemistry and Molecular Biology*, Vol. 27, No. 1, pp. 27-45
36. Obasi N A (2012), "Biochemical Studies on Soil and Air Quality Assessment of Dumpsites on the Enugu-PortHarcourt Expressway, South-East, Nigeria (Ph.D Thesis)", Nigeria: Abia State University Uturu (ABSU).
37. Obute G C, Ndukwu B C and Eze E (2010), "Changes in species diversity and physico-chemical properties of plants in abandoned dumpsites in parts of Por-Harcourt, Nigeria", *Scientia Africana*, Vol. 9, No. 1, pp. 181-193.
38. Ogwueleka T C (2009), "Municipal solid waste characteristics and management in Nigeria", *Iran Journal of Environmental Health Science and Engineering*, Vol. 6, No. 3, pp. 173-180.
39. Okalebo J R, Gathua K W and Woomer P L (1993), *Laboratory methods of soil and plant analysis: A working manual*. Kenya: Marvel EPZ Ltd, Nairobi.
40. Olsen S R and Sommers L E (1982), "Determination of available phosphorus", In: Page F L, Miller R H, Keeney D R (Eds.), *Methods of soil Analysis* (Vol. 2). Madison: American Society Agronomy, pp. 403-07.
41. Osuji C L and Adesiyani S O (2005), "The

- Isiokpo oil pipeline leakage: Total organic carbon/organic matter contents of affected soils”, *Chemical Biodiversity*, Vol. 2, pp. 1079-84.
42. Ozturk M, Ozozen G, Minareci O and Minareci E (2009), “Determination of heavy metals in fish, water and sediments of Avsar dam Lake in Trukey”, *Iran J. Environ. Health Sci. Eng.*, Vol. 6, pp. 73-80.
43. Shanker A K, Cervantes C, Loza-Tavera H and Avadainayagam S (2005), “Chromium toxicity in plants”, *Environment International*, Vol. 31, pp. 739-753.
44. Shauibu U O and Ayodele J T (2002), “Bio-accumulation of four heavy metals in leaves of *Calostropis procera*”, *Journal Chemical Society of Nigeria*, Vol. 27, pp. 26-27.
45. Stumm W and Morgan J J (1981), *Aquatic chemistry: An introduction emphasizing chemical equilibria in natural water* (2nd ed.), New York: John Wiley and Sons.
46. Tessier A, Campbell P G C and Bissom M (1979), “Sequential extraction procedure for the speciation of particulate trace metals”, *Analytical Chemistry*, Vol. 51, No. 7, pp. 844-51.
47. Tokalioglu S, Kantal S and Elci L (2000), “Determination of heavy metals and their speciation in lake sediments by flame atomic absorption spectrophotometer after a four stage sequential extraction procedure”, *Analytical Chemistry Acta*, Vol. 413, pp. 33-40.
48. Uba S, Uzairu A, Harrison G F S, Balarabe M L and Okunola O J (2008), “Assesment of heavy metals bioavailability in dumpsites of Zaira metropolis, Nigeria”, *African Journal of Biotechnology*, Vol. 7(2): 122-130.
49. United State Environmental Protection Agency (USEPA) (1986), Test methods of evaluation of solid waste. In: Contaminated land policies in some industrialized countries. United State: TCB report RO2.
50. Vecera Z, Mikaska P, Zdrahal Z, Docekal B, Buckora M, Tynova Z, Parizek P, Mosna J, Marek J (1999), “Environmental analytical chemistry, Institute of Analytical Chemistry”, *Academy of Sciences of the Zech Republic, Brno. Veveric*, Vol. 97, pp. 61-142.
51. Whitney D A (1999), “Micronutrients: zinc, iron, manganese, and copper”, In: Brown J R (Ed.), Recommended chemical soil test procedures for the north central region. Missouri: Missouri Agric. Experiment Station Bulletin, Vol. 41-44.
52. Yeomans J C and Bremmer J M (1991), “Carbon and Nitrogen analysis of soils by automated combustion techniques”, *Commum Soil Sci. Plant Anal.*, Vol. 22, pp. 843-50.
53. Yoo M S and James B R (2002), “Zinc extractability as a function of pH in organic waste-Contaminated soils”, *Soils Sci.*, Vol. 167, pp. 246-59.
54. Yoon J, Cao X, Zhou Q and Ma L Q (2006), “Accumulation of Pb, Cu and Zn in native plants growing on a contaminated Florida Site”, *Science of the Total Environment*, Vol. 368, pp. 456-464.
55. Zhu Y L, Pilon-Smits E A, Tarun A S, Weber S V, Jouanin L and Terry N (1999), “Cadmium tolerance and accumulation in Indian mustard is enhanced by over expressing gamma-glutamyl cysteine synthetase”, *Plant Physiology*, Vol. 121, pp. 1169-78.



International Journal of Life Sciences Biotechnology and Pharma Research

Hyderabad, INDIA. Ph: +91-09441351700, 09059645577

E-mail: editorijlbpr@gmail.com or editor@ijlbpr.com

Website: www.ijlbpr.com

